

**CERTIFICATE OF ANALYSIS**  
**AICS-0060-027:WTC-mEGFP-MYL2-cl27 (mono-allelic tag)**

<b>Product description</b>	Human iPSC clonal line in which MYL2 has been endogenously tagged with mEGFP using CRISPR/Cas9 technology
<b>Parental cell line</b>	Parental hiPSC line (WTC/AICS-0 at passage 33) derived from fibroblasts reprogrammed using episomal vectors (OCT3/4, shp53, SOX2, KLF4, LMYC, and LIN28). Coriell catalog: GM25256
<b>Publication(s) describing iPSC establishment</b>	Kreitzer et al (2013) Am. J. Stem Cells, 30; 2(2): 119-31
<b>Passage of gene edited iPSC reported at submission</b>	p37 <sup>a</sup>
<b>Number of passages at Coriell</b>	0
<b>Media</b>	mTeSR1
<b>Feeder or matrix substrate</b>	Matrigel
<b>Passage method</b>	Accutase
<b>Thaw</b>	1 million cells (ea vial) in 10 cm plate - ready for passaging in 3-4 days
<b>Seeding density</b>	400K cells/10-cm plate every 4 days or 800K cells/10-cm plate every 3 days (see culture protocol)

<b>Test Description<sup>b</sup></b>	<b>Method</b>	<b>Specification</b>	<b>Result</b>
<b>Post-Thaw Viable Cell Recovery</b>	hiPSC culture on Matrigel	> 50% confluency 3-4 days post-thaw (10 cm plate)	Pass
<b>mEGFP insertion at genomic locus - precise editing</b>	PCR and Sanger sequencing of recombinant and wildtype alleles	C-term insertion of mEGFP in frame with exact predicted recombinant allele junctions. No additional mutations in either allele.	Pass
<b>Copy number</b>	ddPCR <sup>c</sup> assay for mEGFP and RPP30 reference gene <sup>d</sup>	mEGFP/RPP30: ~ 0.5 = Mono-allelic ~ 1.0 = Bi-allelic	Mono-allelic (0.48)
<b>Plasmid integration</b>	ddPCR assay to detect plasmid integration into the genome	AmpR/RPP30: < 0.1 = no plasmid integration	Pass (0.00)
<b>Mutational analysis</b>	Whole exome sequencing <sup>f</sup>	Check for acquired mutations (not detected in p8 <sup>a</sup> parental line) that: 1) Correspond to off-target sites predicted by Cas-OFFinder <sup>e</sup> 2) Affect genes in Cosmic Cancer Gene Census	Sequencing planned
<b>mEGFP localization</b>	Spinning Disk confocal live cell imaging	Localization to sarcomeres in hiPSC-derived cardiomyocytes	Localizes to striations in myofibrils, consistent with localization to thick filaments (myosin-based) within the sarcomere and exclusion from the Z-disk and I-band
<b>Expression of tagged protein</b>	Western blot	Expression of expected size product	Not performed

**CERTIFICATE OF ANALYSIS**  
**AICS-0060-027:WTC-mEGFP-MYL2-cl27 (mono-allelic tag)**

<b>Growth rate</b>	ATP quantitation <sup>g</sup>	Comparable to parental line	Pass (measured at p39) <sup>a</sup>
<b>Expression of stem cell markers</b>	Flow cytometry	Transcription factors: OCT4/SOX2/NANOG $\geq$ 85% Surface markers: SSEA3, TRA-1-60 $\geq$ 85%; SSEA1 $\leq$ 15%	Pass
<b>Germ layer differentiation</b>	Trilineage differentiation <sup>h</sup> as assayed by ddPCR gene expression analysis	Expression of endoderm (SOX17), mesoderm (Brachyury), and ectoderm (PAX6) markers upon directed differentiation to all three germ layers	Pass
<b>Cardiomyocyte differentiation</b>	Modified small molecule differentiation (Lian et al. 2012) <sup>i</sup>	Beating initiated (D7-D14) and Cardiac Troponin T expression (D12-D30) by flow cytometry	Pass
<b>Karyotype</b>	G-banding (30 cell analysis)	Normal karyotype, 46 XY	Pass
<b>Mycoplasma</b>	qPCR (IDEXX)	Negative	Pass
<b>Sterility (bacterial, yeast and fungal testing)</b>	Direct inoculation and incubation for 10 days	No growth after 10 days	Pass
<b>Viral Panel Testing<sup>j</sup></b>	PCR	Negative when assayed for CMV, EBV, HepB, HepC, HIV1, and HPV	Pass
<b>Identity of unedited parental line<sup>k</sup></b>	STR	29 allelic polymorphisms across 15 STR loci compared to donor fibroblasts	Identity matched

<sup>a</sup> This is the number of passages beyond the original parental line (WTC/AICS-0 at passage 33).

<sup>b</sup> All QC assays are performed on stem cells except when noted otherwise.

<sup>c</sup> Droplet digital PCR using Bio-Rad QX200

<sup>d</sup> RPP30 is a reference 2 copy gene used for normalization.

<sup>e</sup> Bae et al (2014) *Bioinformatics*. 30(10): 1473-1475

<sup>f</sup> Nextera rapid capture exome

<sup>g</sup> Promega CellTiter-Glo Luminescent Cell Viability Assay (Catalog #G7571)

<sup>h</sup> STEMCELL Technologies STEMdiff Trilineage Differentiation Kit (Catalog #05230)

<sup>i</sup> Lian et al (2012) *PNAS*. 109(27):E1848-E1857

<sup>j</sup> Viral panel testing was conducted for the parental WTC line prior to editing. Sterility (bacterial, fungal) and mycoplasma testing were conducted in both the parental and edited lines.

<sup>k</sup> STR tests were conducted for the WTC parental line prior to editing. WTC is the only cell line used by AICS. Edited WTC cells were not re-tested because they did not come into contact with any other cell lines.

# CERTIFICATE OF ANALYSIS

## AICS-0060-027:WTC-mEGFP-MYL2-cl27 (mono-allelic tag)

**mEGFP tagging strategy:** Used CRISPR-Cas9 methodology to introduce mEGFP at C-terminus of MYL2 as shown below.




Figure 1: Top: MYL2 locus; Bottom: Zoom in on mEGFP insertion site at MYL2 C-terminus; For more information on tagging transcriptionally silent genes in hiPSCs see: Roberts et al. 2018. Scarless gene tagging of transcriptionally silent genes in hiPSCs to visualize cardiomyocyte sarcomeres in live cells. bioRxiv doi:10.1101/342881

**CERTIFICATE OF ANALYSIS**  
**AICS-0060-027:WTC-mEGFP-MYL2-cl27 (mono-allelic tag)**

---

**Post-thaw imaging:** One vial of distribution lot was thawed (cells were treated with ROCK inhibitor for 24hrs post-thaw - refer to culture protocol). Cultures were observed daily. Colonies were photographed one and three days post-thaw<sup>1,2</sup> using a Leica microscope at 4X and 10x magnification.




Figure 2: Viability and colony formation one day and three days post-thaw

---

<sup>1</sup>Cells may take up to 3 passages to recover after thaw

<sup>2</sup>Morphologies observed post-thaw are representative of cell morphologies observed post-passage

**CERTIFICATE OF ANALYSIS**  
**AICS-0060-027:WTC-mEGFP-MYL2-cl27 (mono-allelic tag)**

---

**Imaging labeled structures in endogenously tagged hiPSC derived cardiomyocytes:** The tagged proteins are expressed endogenously and therefore may not appear as bright as they would in an overexpressed system. For imaging, we plate cells onto high-quality glass bottom 24-well plates (Cellvis) coated with 0.1% w/vol polyethylenimine (PEI) and 25  $\mu\text{g}/\text{ml}$  laminin. Cells are imaged in phenol red-free RPMI 1640 media (Gibco) supplemented with B-27 containing insulin (Gibco). Our most common microscope configuration is a Zeiss spinning disk fluorescence microscope with a Yokogawa CSUX1 head, Hamamatsu CMOS camera, and a 488 laser (GFP). Cells are imaged with a 40x 1.2 NA water immersion objective at 37°C and 5% CO<sub>2</sub> in a temperature-controlled chamber. The approximate laser power measured at the sample for our standard 100x images is  $\sim 2.5$  mW.




Figure 3: Single, mid-level plane of live hiPSC-derived cardiomyocytes expressing mEGFP-tagged MLC-2v protein. Representative high-expressing, relatively mature cell is shown surrounded by more immature cells with lower expression levels (the culture contains cells of varying levels of maturity). Twelve days after the onset of differentiation, cells were plated on PEI and laminin coated glass and imaged in 3D on a spinning disk confocal microscope 47 days later (59 days total after the onset of differentiation). Scale bars, 10  $\mu\text{m}$ .